



The Mathematical Association of Victoria

# Further Mathematics

# 2006 Written Examinations

# Solutions

*These answers and solutions to the 2006 VCE Further Mathematics Written Examinations have been written and published to assist teachers and students in their preparations for future Further Mathematics Examinations. They have been published without the relevant questions to avoid any breaches of copyright. They are suggested answers and solutions only and do not necessarily reflect the views of the Victorian Curriculum and Assessment Authority Assessing Panels.*

© The Mathematical Association of Victoria 2007

These answers and solutions are licensed to the purchasing school or educational organisation with permission for copying within that school or educational organisation. No part of this publication may be reproduced, transmitted or distributed, in any form or by any means, outside purchasing schools or educational organisations or by individual purchasers without permission.

**Published by The Mathematical Association of Victoria**  
"Cliveden", 61 Blyth Street, Brunswick, 3056  
Phone: (03) 9380 2399 Fax: (03) 9389 0399  
E-mail: [office@mav.vic.edu.au](mailto:office@mav.vic.edu.au) website: <http://www.mav.vic.edu.au>

---

**2006 Further Mathematics**  
**Written Examination 1 (Facts, skills and applications)**  
**Suggested answers and solutions**

**SECTION A (Multiple-Choice)****Answers****Core**

1. **D**   2. **B**   3. **E**   4. **C**   5. **D**  
 6. **B**   7. **A**   8. **C**   9. **A**   10. **C**  
 11. **D**   12. **C**   13. **A**

**SECTION B (Multiple-Choice)****Answers****Module 1 Number patterns and applications**

1. **D**   2. **C**   3. **C**   4. **A**   5. **D**  
 6. **E**   7. **B**   8. **D**   9. **E**

**Module 2 Geometry and trigonometry**

1. **B**   2. **D**   3. **A**   4. **C**   5. **C**  
 6. **D**   7. **E**   8. **A**   9. **D**

**Module 3 Graphs and relations**

1. **D**   2. **E**   3. **C**   4. **A**   5. **B**  
 6. **C**   7. **E**   8. **A**   9. **C**

**Module 4 Business-related mathematics**

1. **B**   2. **C**   3. **E**   4. **B**   5. **E**  
 6. **C**   7. **D**   8. **A**   9. **B**

**Module 5 Networks and decision mathematics**

1. **B**   2. **B**   3. **D**   4. **D**   5. **E**  
 6. **C**   7. **C**   8. **D**   9. **D**

**Module 6 Matrices**

1. **B**   2. **D**   3. **A**   4. **B**   5. **D**  
 6. **E**   7. **C**   8. **E**   9. **B**

**Core****Question 1 [D]**

Temperature ( $^{\circ}$  Celsius) and Town (Beachside and Flattown) are numerical and categorical variables respectively.


**Question 2 [B]**

$$\begin{aligned}\text{Range} &= \text{Maximum} - \text{Minimum} \\ &= 38 - 15 \\ &= 23\end{aligned}$$

**Question 3 [E]**

Flattown	89	89	334	55677788	0012	56	
	1	2	2	3	3	4	4


The distribution of maximum temperatures for Flattown is best described as approximately symmetric with outliers.

**Question 4 [C]**

The shaded area less than 48 cm is approximately 16%.

The number of boys with head circumference of less than 48.0 cm is closest to  $16\% \times 400 = 64$

### Question 5 [D]



The % of students who passed the test (ie 30 or above) is

$$7 + 11 + 14 + 16 + 18 + 12 = 78\%$$

### Question 6 [B]

The median point lies at 50%.

The % of students from 0 to 30-35 is

$$2 + 3 + 2 + 3 + 6 + 6 + 7 + 11 = 40$$

The % of students from 0 to 40-45 is

$$2 + 3 + 2 + 3 + 6 + 6 + 7 + 11 + 14 = 54$$

So the median lies between 40-45.

### Question 7 [A]

$$r = -0.5675, \bar{x} = 4.56, s_x = 2.61,$$

$$\bar{y} = 23.93 \text{ and } s_y = 6.98$$

$$b = r \frac{s_y}{s_x} = -0.5675 \left( \frac{6.98}{2.61} \right) \approx -1.52$$

$$a = \bar{y} - b\bar{x} = 23.93 + 1.52(4.56) \approx 30.9$$

$$y = 30.9 - 1.52x$$

### Question 8 [C]

**Actual**

When Waist = 80 cm, weight = 67 kg

**Estimate**

$$\text{Weight} = -20 + (1.11 \times 80) = 68.8$$

**Residual =**

$$y_{act} - y_{est} = 67 - 68.8 = -1.8 \approx -2 \text{ kg}$$


### Question 9 [A]

Input data into 2 lists.

$x$  into  $L_1$  and  $y$  into  $L_2$ .  $L_3 = x^2$

L1	L2	L3	3
0	5	0	
1	7	1	
2	14	4	
3	33	9	
4	58	16	
5	76	25	
6	124	36	

$$L3(1)=0$$



To find the least squares regression line for the transformed data:

STAT

CALC

8: LinReg (a+bx)  $L_3, L_2$

LinReg

y=a+bx

a=7.147035573

b=2.938700506

r<sup>2</sup>=.991697752

r=.9958402241



So the equation is  $y = 7.1 + 2.9x^2$

### Question 10 [C]

Mar	April	May
35	99	75

Three mean moving average for April

$$= \frac{35 + 99 + 75}{3} = 69.66... \approx 70$$

**Question 11** [D]

The seasonal indices add to 12.

The seasonal index for October

$$= 12 - (1.30 + 1.21 + 1.00 + 0.95 + 0.95$$

$$+ 0.86 + 0.86 + 0.89 + 0.94 + 0.99 + 1.07))$$

$$= 0.98$$

**Question 12** [C]

$$\text{Deseasonalised} = \frac{\text{Actual}}{\text{Seasonal Index}}$$

$$= \frac{330}{0.94} = 351$$

**Question 13** [A]

Deseasonalised :

$$= 373.3 - (3.38 \times 6)$$

$$= 353.0$$

Actual is Deseasonalised  $\times$  Seasonal Index

$$= 353.02 \times 0.86$$

$$\approx 304$$

**Module 1 Number patterns****Question 1** [D]

An arithmetic sequence has a common difference  $d$ .

For  $-4, -1, 2, 5, 8, \dots$ ,

$$d = -1 - (-4) = 3$$

**Question 2** [C]

The first 3 terms of a geometric sequence are 6,  $x$ , 54.

$$\frac{x}{6} = \frac{54}{x}$$

$$x^2 = 324$$

$$x = 18$$

**Question 3** [C]

At the start of the second year, the

farmer has  $(50 \times 1.84) - 40 = 52$  sheep

**Question 4** [A]

$$S_1 = 50, d = -40, r = 1.84$$

The difference equation which models this growth of sheep over time is:

$$S_{n+1} = 1.84S_n - 40 \text{ where } S_1 = 50$$

**Question 5** [D]

For  $f_{n+1} - f_n = 5$  where  $f_1 = -1$

$$f_2 = 5 + f_1 = 5 - 1 = 4$$


$$f_3 = 5 + f_2 = 5 + 4 = 9$$

So the sequence is  $-1, 4, 9, \dots$

**Question 6** [E]

$$a = 12, r = 1.03$$

$$t_{15} = 12 \times 1.03^{14} = 18.15 \approx 18.2$$

**Question 7** [B]

As the graph is non linear, the difference equation is **not** arithmetic (A, C and D)

The sequence is 4, 9, 19, 39, 79

So the difference equation is

$$t_{n+1} = 2t_n + 1, t_1 = 4$$

**Question 8** [D]

$$t_n = t_{n-1} + t_{n-2} \text{ where } t_1 = 1 \text{ and } t_2 = 2$$

$$t_3 = t_2 + t_1 = 2 + 1 = 3$$

$$t_4 = t_3 + t_2 = 3 + 2 = 5$$

$$t_5 = t_4 + t_3 = 5 + 3 = 8$$

So the total number of stamps after 5

$$\text{weeks} = 1 + 2 + 3 + 5 + 8 = 19$$

**Question 9** [E]

$$a = 73.4, r = \frac{380}{400} = \frac{361}{380} = 0.95$$


$$S_{\infty} = \frac{400}{1 - 0.95} = 8000g = 8 \text{ kg}$$

Roh's eventual body weight will be

$$73.4 + 8 = 81.4 \text{ kg}$$

## Module 2 Geometry and trigonometry


### Question 1 [B]



$$\sin \theta = \frac{142}{215}$$

$$\theta = \sin^{-1} \frac{142}{215} \approx 41^\circ$$

### Question 2 [D]



$$\frac{45}{\sin 54^\circ} = \frac{RT}{\sin 108^\circ}$$


$$RT = \frac{45 \sin 108^\circ}{\sin 54^\circ} = 52.9^\circ \approx 53^\circ$$

### Question 3 [A]



The difference in height between A and B is  $300 - 200 = 100$  m

### Question 4 [B]



$$V = (8 \times 4 \times 24) + (0.5 \times 2 \times 8 \times 24) = 960 \text{ m}^3$$

### Question 5 [E]


Using Heron's Formula:

$$S = \frac{1}{2}(a + b + c) = \frac{1}{2}(36 + 58 + 42) = 68$$

$$A = \sqrt{S(S-a)(S-b)(S-c)}$$

$$= \sqrt{68(68-36)(68-58)(68-42)}$$

### Question 6 [D]



$$V = l \times w \times h$$

$$6 = 1.5 \times 2 \times h,$$

$$\text{So } h = 2$$

$$TSA = 2(1.5 \times 2) + 2(2 \times 2) + 2(1.5 \times 2)$$

$$= 6 + 8 + 6$$

$$= 20 \text{ m}^2$$


### Question 7 [E]

Using Similar Triangles  $\triangle CAB$  and  $\triangle EDB$

$$\frac{24}{36} = \frac{DE}{27}$$

$$DE = \frac{24 \times 27}{36} = 18 \text{ cm}$$

### Question 8 [A]




$$\begin{aligned} x^2 &= 50^2 - 40^2 \\ &= 2500 - 1600 \\ &= 900 \end{aligned}$$

$$x = 30$$

$$d = 50 - 30 = 20 \text{ cm}$$


### Question 9 [D]



From the diagram, the bearing of P from M is between  $180^\circ$  and  $270^\circ$


### Module 3 Graphs and relations

#### Question 1 [D]




The equation of the line passing through the point  $(1, 2)$  is  $y = 2$

#### Question 2 [E]



From the graph, there are 4 times when the depth of the water is 10m.

#### Question 3 [C]



From the graph it can be seen that the lamp runs out of gas after 4.5 hours.



**Question 4 [A]**

The gradient for the straight line joining  $(0, 332.5)$  to  $(4.5, 220)$  is:

$$\frac{220 - 332.5}{4.5} = -25$$

The  $M$ -intercept is 332.5

The gradient from  $(4.5, 220)$  onwards is 0.

The rule would be

$$M = \begin{cases} 332.5 - 25t & \text{for } 0 \leq t \leq 4.5 \\ 220 & \text{for } 4.5 < t \leq 6 \end{cases}$$

**Question 5 [B]**


The equation  $12x - 4y = 0$  can be rewritten as  $y = 3x$ .

So the line **does not** have a slope of 12.

**Question 6 [C]**

The point of intersection of two lines is  $(2, -2)$  so  $(2, -2)$  must be a point on both lines.

Substituting  $(2, -2)$  into  $2x + 2y = 0$  gives  $4 - 4 = 0$  which is true. So one of these two lines could be  $2x + 2y = 0$ .

**Question 7 [E]**

One of the constraints defining the feasible region indicates that the total number of cows and sheep cannot exceed 80.

**Question 8 [A]**

The cost equation is  $C = 400 + 50x$ .


Breakeven occurs when Cost = Sell

For 10 frying pans the cost and selling prices are:

$$\text{Cost} = 400 + 50 \times 10 = 900$$

$$\text{Sell} = 90 \times 10 = 900$$

So the manufacturer could break even by selling 10 frying pans at \$90 each.

**Question 9 [C]**

$(6, 2)$  lies within the feasible region.

**Module 4 Business-related mathematics****Question 1 [B]**

$$I = \frac{Pr t}{100} = \frac{4000 \times 5 \times 1}{100} = 200$$

\$200 interest is earned in the first year.

**Question 2 [C]**

Minimum balance for October is \$473.92.

$$I = \frac{Pr t}{100} = \frac{473.92 \times 0.15 \times 1}{100 \times 1} = \$0.71$$

Interest paid for October is \$0.71.

**Question 3 [E]**

A perpetuity is an annuity where a permanently invested sum of money provides regular payments that continue forever.

Yearly interest =  $\$584 \times 12 = \$7008$

$$P = \frac{100I}{rt} = \frac{100 \times 7008}{6.2 \times 1} \approx \$113000$$

Grandpa needs to invest \$113000 in the perpetuity.

**Question 4 [B]**

$$1.1x = 825$$

$$x = \frac{825}{1.1} = 750 \quad (\text{without GST})$$

$$\text{GST} = 825 - 750 = \$75.00$$

**Question 5 [E]**

Total Number of copies

$$= \frac{48000 - 21000}{0.04}$$

$$= 675000$$

**Question 6 [C]**

$$\text{Principal} = \$2000 - \$200(\text{deposit}) = \$1800$$

$$\text{Total Instalments} = 36 \times 68 = \$2448$$

$$I = 2448 - 1800 = 648$$

$$r_f = \frac{100 \times 648}{1800 \times 3} = 12\%$$


Annual flat rate of interest is 12%.

**Question 7 [D]**

Price paid for lawn mower = \$368

$$\$368 + \$80 (\text{Trade in}) = \$448$$

$$\frac{448}{0.8} = \$560.00 \quad (\text{Original cost})$$

**Question 8 [A]**

The investment has interest compounding annually and is credited annually.

**Question 9 [B]**

To calculate the monthly repayment

```

N=60
I%=9.2
PV=18000
PMT=-375.40000...
FV=0
P/Y=12
C/Y=12
PMT:[FV] BEGIN
  
```

After the tenth repayment

```

N=10
I%=9.2
PV=18000
PMT=-375.40000...
FV=15542.39977
P/Y=12
C/Y=12
PMT:[FV] BEGIN
  
```


$$\$18000 - 15542.40 = \$2457.60$$

Jenny has paid \$2457.60 of the principal immediately following the tenth repayment.




## Module 5 Networks and decision mathematics

### Question 1 [B]




The number of vertices with an odd degree is 2.

### Question 2 [B]




All intersections can be reached from intersection 5.

### Question 3 [D]



As there are two vertices with odd degree, at least two Eulerian paths exist.

### Question 4 [D]




The minimal spanning tree for the network will include the edge that has the weight of 9.


### Question 5 [E]

Activity	Immediate predecessors
A	—
B	—
C	—
D	A
E	B
F	D, E
G	C
H	C
I	F, G
J	H, I

The directed graph which represents this project is:



### Question 6 [C]




The capacity of the cut is  $e + c + b + a$ .  
Note:  $d$  is heading back into the cut towards the source, so is not included.

### Question 7 [C]

A six-team basketball competition where all teams play each other once would represent a complete graph with six vertices.


### Question 8 [D]

Euler's formula applies to planar graphs which have no edges crossing.



This diagram cannot be drawn without edges crossing.

### Question 9 [D]



If activity B is reduced by 4 hours, the project completion time is also reduced by 4 hours.

## Module 6 Matrices

### Question 1 [B]

The matrix  $\begin{bmatrix} 12 & 36 \\ 0 & 24 \end{bmatrix} = 12 \begin{bmatrix} 1 & 3 \\ 0 & 2 \end{bmatrix}$

### Question 2 [D]

$A = \begin{bmatrix} -2 \\ 0 \end{bmatrix}$ ,  $B = \begin{bmatrix} 0 & 9 \end{bmatrix}$ ,  $C = \begin{bmatrix} 2 \end{bmatrix}$

The order of B is  $1 \times 2$ ; the order of C is  $1 \times 1$ . If BC is defined then the number of columns of B must equal the number of rows of C.

Order BC =  $(1 \times 2) \times (1 \times 1)$  ... this is not so and therefore the matrix BC is not defined.

**Question 3 [A]**

$$A = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, B = \begin{bmatrix} 2 & 1 \\ 1 & 0 \end{bmatrix}, C = \begin{bmatrix} 1 & -1 \\ -1 & 1 \end{bmatrix}$$

Now  $A$  is the  $2 \times 2$  unit matrix and so

$$A^3 \text{ is also } \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$B - C = \begin{bmatrix} 2 & 1 \\ 1 & 0 \end{bmatrix} - \begin{bmatrix} 1 & -1 \\ -1 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 2 \\ 2 & -1 \end{bmatrix}$$

$$\begin{aligned} A^3(B - C) &= \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 2 & -1 \end{bmatrix} \\ &= \begin{bmatrix} 1 & 2 \\ 2 & -1 \end{bmatrix} \end{aligned}$$

**Question 4 [B]**

	Athletics	Cross country	Swimming
2004	Green	Green	Blue
2005	Green	Red	Blue
2006	Blue	Green	Blue

The total number of competitions won by each if the three teams in each of these three years is

$$\begin{array}{c} B \quad G \quad R \\ \begin{array}{l} 2004 \begin{bmatrix} 1 & 2 & 0 \end{bmatrix} \\ 2005 \begin{bmatrix} 1 & 1 & 1 \end{bmatrix} \\ 2006 \begin{bmatrix} 2 & 1 & 0 \end{bmatrix} \end{array} \end{array}$$

**Question 5 [D]**

The new price matrix

$$MP = \begin{bmatrix} 1.2 & 0 \\ 0 & 1.35 \end{bmatrix}$$

$$\begin{bmatrix} 145 & 210 & 350 \\ 185 & 270 & 410 \end{bmatrix}$$

$$= \begin{bmatrix} 1.2 \times 145 + 0 & 1.2 \times 210 + 0 & 1.2 \times 350 + 0 \\ 0 + 1.35 \times 185 & 0 + 1.35 \times 270 & 0 + 1.35 \times 410 \end{bmatrix}$$

$$= \begin{bmatrix} 174 & 252 & 420 \\ 249.75 & 364.50 & 553.50 \end{bmatrix}$$

**Question 6 [E]**

$$XA = X \begin{bmatrix} 1 & 3 \\ 6 & 4 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} 4 & 1 \\ 1 & 4 \\ 3 & 5 \end{bmatrix}$$

Let the order of  $X$  be  $m \times n$ . The order of  $A$  is  $3 \times 2$ .

$$\begin{array}{c} \text{Order } XA = (m \times n) \times (3 \times 2) \\ \quad \quad \quad \uparrow \quad \quad \quad \uparrow \\ \quad \quad \quad (m \times 2) \end{array}$$

For this to exist then  $n = 3$ .

But  $(m \times 2) = (3 \times 2)$  and so  $m = 3$  as well.

**Question 7 [C]**

For simultaneous equations to have a unique solution, the determinant cannot be zero.

$4x + 2y = 10$	$x = 0$	$x - y = 3$	$2x + y = 5$	$x = 8$
$2x + y = 5$	$x + y = 6$	$x + y = 3$	$2x + y = 10$	$y = 2$
(1)	(2)	(3)	(4)	(5)

$$1. \det(1) = \begin{vmatrix} 4 & 2 \\ 2 & 1 \end{vmatrix} = 4 - 4 = 0$$

$$2. \det(2) = \begin{vmatrix} 1 & 0 \\ 1 & 1 \end{vmatrix} = 1 - 0 = 1$$

$$3. \det(3) = \begin{vmatrix} 1 & -1 \\ 1 & 1 \end{vmatrix} = 1 - (-1) = 2$$

$$4. \det(4) = \begin{vmatrix} 2 & 1 \\ 2 & 1 \end{vmatrix} = 2 - 2 = 0$$

$$5. \det(5) = \begin{vmatrix} 1 & 0 \\ 0 & 1 \end{vmatrix} = 1 - 0 = 1$$

There are 3 sets of simultaneous equations with a unique solution.

**Question 8 [E]**

95% of those who had their last holiday in Australia would have their next holiday in Australia.

20% of those who had their last holiday in Australia would have their next holiday overseas.

The transition matrix for this situation

is  $\begin{bmatrix} 0.95 & 0.80 \\ 0.05 & 0.20 \end{bmatrix}$

**Question 9 [B]**

The movement of birds is described by the transition matrix:

$$\begin{array}{cc} & \begin{array}{cc} A & B \end{array} \\ \begin{array}{c} A \\ B \end{array} & \begin{bmatrix} 0.8 & 0 \\ 0.2 & 1 \end{bmatrix} \end{array}$$

In the long term, the number of birds that settle at location A will gradually decrease to zero. This can be confirmed by multiplying this matrix by itself over and over again.

**2006 Further Mathematics  
Written Examination 2 (Analysis task)  
Suggested answers and solutions**

**Core**

**Question 1**

**a 18-month-old boys**

Input Column 1 from Table 1 into calculator:

STAT

CALC


Enter 1 : 1-Var Stats for your list.

```
1-Var Stats
 $\bar{x}$ =82.67142857
 $\Sigma x$ =1157.4
 $\Sigma x^2$ =95875.76
 $Sx$ =3.841559894
 $\sigma x$ =3.70181974
 $\downarrow n$ =14
```

The standard deviation ( $S_x$ ) for 18 months is 3.8 (correct to one decimal place).

**b**  $Z\text{-score} = \frac{x - \bar{x}}{s_x} = \frac{83.1 - 89.3}{4.5} = -1.4$  (correct to one decimal place)

**c.** A 36-month-old boy has a standardised height  $Z$  value of 2.




Approximately 97.5% of 36-month-old boys will be shorter than this child.

**d. 18-month-old boys five number summary**


```
1-Var Stats
 $\uparrow n$ =14
minX=76
Q1=80
Med=83
Q3=85.8
maxX=89.8
```





- e. From the boxplot for 27-month-old boys, the median height is 89.5cm.
- f. The median of heights for 18-month-old, 27-month-old and 36-month-old boys is 83cm, 89.5cm and 95cm respectively. The median is increasing, suggesting that height and age are positively related.

## Question 2




The least squares regression line is  $\text{Height} = 75.4 + 0.53 \times \text{age}$

The correlation coefficient is  $r = 0.7541$

- a. On average, the height of a boy increases by **0.53** cm for each one-month increase in age.
- b. i The coefficient of determination  $r^2 = (0.7541)^2 = 0.5687 = 56.9\%$  (correct to one decimal place)
- ii 56.9% of the variation in height can be explained by the variation in age.

### Question 3

a.



b. Input data into 2 lists  $L_1, L_2$  and do a 3 **Med-Med**. Line

**Med-Med**

$$y = ax + b$$

$$a = .6666666667$$

$$b = 70.66666667$$

The three median line is: height =  $70.7 + 0.7 \times \text{age}$  (correct to 1 decimal place)

c. As there is an outlier at (20, 93), the three median line is the preferable model to use as the least squares regression line is significantly affected by outliers.

### Module 1: Number patterns and applications

#### Question 1

a. At the end of the second day there are  $48000 - 2(3000) = 42000$  kg of fruit

b. The value of  $d$  is  $-3000$ .

c. If all the fruit is picked from the trees then:  $-3000 \times n + 48000 = 0$   
So  $n = 16$  days.

#### Question 2

a.  $r = \frac{t_2}{t_1} = \frac{500}{625} = 0.8$  and  $r = \frac{t_3}{t_2} = \frac{400}{500} = 0.8$

So  $r = 0.8$ , as required.

b.  $a = 625$   $r = 0.8$  and  $t_n = ar^{n-1}$

$$t_5 = ar^{5-1} = 625 \times 0.8^4 = 256$$

The gardeners will work 256 hours in the fifth month.

c. The number of hours that the gardeners will work in the  $n$ th month after planting is

$$H_n = 625 \times 0.8^{n-1}$$

d.  $t_6 = 204.8$  hours and  $t_7 = 163.84$  hours

$$204.8 - 163.84 = 40.96$$

The gardeners work 41 hours more (to the nearest hour).

e. Solve  $100 = 625 \times 0.8^{n-1}$ , using the graphics calculator.



Therefore the gardeners work less than 100 hours during the 10<sup>th</sup> month.

f. In the next nine months, i.e. months 3 to 12, the gardeners work  $S_{12} - S_3$ .

$$= 2910.25 - 1525$$

$$= 1385.25$$

Answer: 1385 hours (to the nearest hour).

### Question 3

a. The volume of water,  $V_n$ , in the tank on the morning of the  $n$ th day is modelled by the difference equation  $V_{n+1} = rV_n + d$  where  $V_1 = 45000$  litres

As 10% of the volume is used,  $r = 0.9$

As 2000 litres is added to the tank,  $d = 2000$

b.  $V_{n+1} = 0.9V_n + 2000$

$$V_1 = 45000 \text{ litres}$$

$$V_2 = 0.9V_1 + 2000 = 0.9(45000) + 2000 = 42500$$

$$V_3 = 0.9V_2 + 2000 = 0.9(42500) + 2000 = 40250$$

$$V_4 = 0.9V_3 + 2000 = 0.9(40250) + 2000 = 38225$$

There is 38225 litres of water in the tank on the morning of the fourth day.

- c. To find when the tank will first be below 30000 litres, solve the difference equation:

```

Plot1 Plot2 Plot3
nMin=1
u(n)=0.9u(n-1)+
2000
u(nMin)=45000
v(n)=
v(nMin)=
w(n)=

```

$n$	$u(n)$
1	38225
2	36403
3	34762
4	33286
5	31957
6	30762
7	29686
8	
9	
10	

$n=10$


The tank will first be below 30000 litres during the 10<sup>th</sup> day.

- d. There will be 20000 litres in the tank each morning. It drops 10% to 18000 litres during the afternoon but then 2000 litres is added each evening bringing it back to 20000.

## Module 2: Geometry and trigonometry

### Question 1

A farmer owns a flat allotment of land in the shape of triangle  $ABC$  shown below.  
 Boundary  $AB$  is 251 metres.  
 Boundary  $AC$  is 142 metres.  
 Angle  $BAC$  is  $45^\circ$ .



a.  $\angle AXY = 180^\circ - (45^\circ + 90^\circ) = 45^\circ$

b.  $\cos 45^\circ = \frac{55}{AX}$

$$AX = \frac{55}{\cos 45^\circ}$$

$$= 77.8 \quad (\text{correct to 1 decimal place})$$

c. From the diagram above, the bearing of B from A is  $78^\circ - 45^\circ = 033^\circ T$

d.  $\tan 45^\circ = \frac{XY}{55}$

$$XY = 55 \tan 45^\circ$$

$$= 55$$

$$XC = \sqrt{55^2 + 87^2} = 102.9 \text{ (correct to 1 decimal place)}$$

e.

$$\text{Area of } \triangle ABC = \frac{1}{2} \times 251 \times 142 \times \sin 45^\circ$$

$$= 12601.34$$

$$\approx 12601 \text{ m}^2 \text{ (correct to nearest square metre)}$$

*Alternatively*, Heron's Rule can be used to find the area of the triangle.

Using the Cosine rule,  $BC = \sqrt{251^2 + 142^2 - (2 \times 251 \times 142 \cos 45^\circ)} = 181 \text{ m}$

$$s = \frac{1}{2}(a + b + c)$$

$$= \frac{1}{2}(251 + 142 + 181)$$

$$= 287$$

$$\text{Area} = \sqrt{s(s-a)(s-b)(s-c)}$$

$$= \sqrt{287(287-251)(287-142)(287-181)}$$

$$= 12601.699$$

$$\approx 12602 \text{ m}^2 \text{ (correct to the nearest square metre)}$$

f. i. Using the Cosine rule,  $BC = \sqrt{251^2 + 142^2 - (2 \times 251 \times 142 \cos 45^\circ)}$

$$= \sqrt{32759}$$

$$= 180.996$$

$$= 181 \text{ m (correct to 1 decimal place).}$$

ii. Using the Sine rule,  $\frac{181}{\sin 45^\circ} = \frac{142}{\sin \theta}$


$$\sin \theta = \frac{142 \sin 45^\circ}{181}$$

$$\theta = \sin^{-1}\left(\frac{142 \sin 45^\circ}{181}\right)$$

$$\theta = 33.7^\circ \text{ (correct to 1 decimal place)}$$



**g.**




As the  $\triangle AMN$  is isosceles, solve:

$$3200 = \frac{1}{2} MN^2$$

$$6400 = MN^2 \text{ and so } MN \text{ is } 80\text{m.}$$

## Question 2

**a.**




$$\tan 20^\circ = \frac{PQ}{125} \text{ and so } PQ = 125 \tan 20^\circ$$

$$PQ = 45.5 \text{ m (correct to 1 decimal place)}$$

**b.** The angle of depression is given by  $\theta$  where  $\tan \theta = \frac{45.5}{98}$ .

$$\theta = \tan^{-1} \frac{45.5}{98} = 24.9^\circ \text{ (correct to 1 decimal place)}$$

### Question 3




a. From the diagram,  $r = \frac{3}{2} = 1.5$  m

b.

$$\begin{aligned}
 V_{\text{cylinder}} &= \pi r^2 h \\
 &= \pi \times 1.5^2 \times (2.4 - (2 \times 0.25)) \\
 &= 13.43 \\
 &\approx 13 \text{ m}^3 \text{ (correct to the nearest cubic metre)}
 \end{aligned}$$

### Module 3: Graphs and relations

#### Question 1




a. i. The call-out fee to travel a distance of 20 km is \$30.

ii. The maximum distance travelled for a call-out fee of \$10 is 5 km.

b. Refer to the diagram above for a call-out fee of \$50 being charged to travel distances of more than 30 km but less than or equal to 40 km.

## Question 2



- a. Two coordinates on the line are (0,50) and (160,30)

$$\text{Gradient } m = \frac{30 - 50}{160 - 0} = -\frac{1}{8}$$


$$\text{Equation of the line is } y = -\frac{1}{8}x + 50$$

- b. Solving  $-\frac{1}{8}x + 50 = 0$ , gives  $x = 400$

He had already travelled 160 km and so  $400 - 160 = 240$  km is the distance that he needs to travel further before the tank is empty.

- c. When the tank is completely full it holds  $12 + (18 \times 3.5) = 75$  litres.

## Question 3



(Feasible region shaded)

- a. For  $20x + 25y = 200$ ,

The  $x$ -intercept:  $20x = 200$  so  $x = 10$  (10, 0)

The  $y$ -intercept:  $25y = 200$  so  $y = 8$  (0, 8)

Refer to diagram above for the line.

- b. In any one day, the number of dogs clipped ( $y$ ) is at least twice the number of dogs washed ( $x$ ).

Inequality 4 is  $y \geq 2x$

- c. i. Refer to the diagram above for the boundaries of the region represented by Inequalities 1 to 4.

- ii. From the graph, when  $y = 5$ ,  $x = 2.5$ .

So the maximum number of dogs that could be washed is 2.

- d. The profit from washing one dog is \$40 and the profit from clipping one dog is \$30 so the total profit is given by the equation:  $P = 40x + 30y$

- e. i. Using the profit equation  $P = 40x + 30y$  and the feasible region coordinates:

$$(0, 8) \quad P = 0 \times 40 + 8 \times 30 = \$240$$

$$(2, 4) \quad P = 2 \times 40 + 4 \times 30 = \$200$$

$$(2, 5) \quad P = 2 \times 40 + 5 \times 30 = \$230$$

$$(2, 6) \quad P = 2 \times 40 + 6 \times 30 = \$260$$

Maximum total profit is achieved if 2 dogs are washed and 6 dogs are clipped.

- ii. Maximum total profit is \$260.

## Module 4: Business-related mathematics

### Question 1

- a. i. Annual depreciation =  $60000 \times 0.1 = \$6000$

- ii. After three years, the value of the machine will be  $60000 - (6000 \times 3) = \$42000$

- iii.  $12000 = 60000 - 6000n$

$$6000n = 48000 \quad \text{so } n = 8$$

The machine is worth \$12000 after 8 years.


- b.  $V = 60000 \times (0.85)^n$ .

- i. Annual depreciation =  $1 - 0.85 = 0.15 = 15\%$

- ii. When  $n = 3$ ,  $V = 60000 \times 0.85^3 = \$36847.50$ .


The machine is worth \$36847.50

- iii. Solve  $12000 = 60000 \times 0.85^n$



The machine will first fall below \$12000 at the end of 10 years.

c. Solve  $60000 - 6000n = 60000 \times 0.85^n$



The machine will be less using Flat Rate rather than Reducing Balance Depreciation at the end of 7 years.

## Question 2

Cost of a new machine =  $60000 \times (1.02)^8 = \$70300$  (correct to the nearest dollar)

## Question 3

a.  $P = \$7000$ ,  $r = 6.25\%$ ,  $t = 8$

$$I = \frac{7000 \times 6.25 \times 8}{100} = \$3500$$

Investment = Principal + Interest =  $\$7000 + \$3500 = \$10500$

b.  $P = \$10000$ ,  $r = 6\%$  p.a.,  $t = 8$  years but compounded **quarterly**.

$$A = 10000 \times \left(1 + \frac{6}{400}\right)^{32}$$

$$= \$16103.24$$

$$\approx \$16103$$

After 8 years the investment is worth \$16103 (correct to the nearest dollar)

c.

```
N=96
I%=6.5
PV=500
PMT=200
FV=-25935.30411
P/Y=12
C/Y=12
PMT:[END] BEGIN
```

After 8 years, the investment is worth \$25935 (correct to the nearest dollar)

## Question 4

```
N=24
I%=10
PV=20000
PMT=-922.89852...
FV=0
P/Y=12
C/Y=12
PMT:[END] BEGIN
```

The monthly repayment is \$922.90.


The total amount of interest:

$$= (24 \times 922.90) - 20000 = \$2150 \text{ (correct to the nearest dollar)}$$



## Module 5: Networks and decision mathematics

### Question 1



a George must play the guitar.

b.

Person	Position
Harriet	Drums
Ian	Saxophone
Keith	Keyboards

### Question 2


a.

Matrix 1

		loser				
		G	H	I	J	K
winner	G	<b>0</b>	0	1	0	1
	H	1	<b>0</b>	1	1	0
	I	0	0	<b>0</b>	1	0
	J	1	0	0	<b>0</b>	0
	K	0	1	1	1	<b>0</b>

The figures in bold in Matrix 1 are all zero as no musician can compete against his/herself.

b.



Missing edge – Josie defeats George.

c. Matrix 2 shows the 2 step dominances.

**Matrix 2**

	<i>G</i>	<i>H</i>	<i>I</i>	<i>J</i>	<i>K</i>
<i>G</i>	0	1	1	2	0
<i>H</i>	1	0	1	1	1
<i>I</i>	1	0	0	0	0
<i>J</i>	0	0	1	0	1
<i>K</i>	2	0	1	$x$	0

George defeated Keith and Keith defeated Ian so George has a 2 step dominance over Ian.

d.  $x = 2$

e.


Musician	Dominance value (one step)	Dominance value (two step)	Total
George	2	4	6
Harriet	3	4	7
Ian	1	1	2
Josie	1	2	3
Keith	3	5	8

Keith came first and Ian came last.

### Question 3

Activity	Immediate predecessors
<i>A</i>	—
<i>B</i>	—
<i>C</i>	—
<i>D</i>	<i>A</i>
<i>E</i>	<i>B</i>
<i>F</i>	<i>C</i>
<i>G</i>	<i>D, E</i>
<i>H</i>	<i>F</i>
<i>I</i>	<i>G, H</i>

The network for the activities above is:



- b. There are 5 non-critical activities: A, C, D, F and H.
- c. The critical path for the project is B E G I.
- d. Duration of activity I =  $19 - 12 = 7$  hours
- e. C, F and H cannot add to more than the latest starting time for activity I which is 12. Time for F&H =  $12 - C = 12 - 3 = 9$  hours. As activity C has a float time of 1 hour, the time for F&H can be no more than 8 hours.

## Module 6: Matrices

### Question 1

- a. The order of Q is  $2 \times 3$ .

b. i.  $M = QP = \begin{bmatrix} 2500 & 3400 & 1890 \\ 1765 & 4588 & 2456 \end{bmatrix} \begin{bmatrix} 14.50 \\ 21.60 \\ 19.20 \end{bmatrix} = \begin{bmatrix} 145978 \\ 171848.50 \end{bmatrix}$

- ii. Matrix M gives the total revenue from selling products A, B and C at Eastown and Noxland.

c. Order of  $\begin{bmatrix} 14.50 \\ 21.60 \\ 19.20 \end{bmatrix}$  is  $(3 \times 2)$  and order of  $\begin{bmatrix} 2500 & 3400 & 1890 \\ 1765 & 4588 & 2456 \end{bmatrix}$  is  $(2 \times 3)$

PQ is not defined as the number of columns of P  $\neq$  number of rows in Q.

### Question 2

- a. The transition matrix T is:

$$T = \begin{array}{ccc|c} \text{This week} & & & \\ S & E & N & \\ \hline \begin{bmatrix} 0.80 & 0.09 & 0.10 \\ 0.12 & 0.76 & 0.05 \\ 0.08 & 0.15 & 0.85 \end{bmatrix} & S & E & \text{Next week} \\ & N & & \end{array}$$

b.  $K_0 = \begin{bmatrix} 300000 \\ 120000 \\ 180000 \end{bmatrix}$

c.  $K_1 = T \times K_0 = \begin{bmatrix} 0.80 & 0.09 & 0.10 \\ 0.12 & 0.76 & 0.05 \\ 0.08 & 0.15 & 0.85 \end{bmatrix} \begin{bmatrix} 300000 \\ 120000 \\ 180000 \end{bmatrix} = \begin{bmatrix} 268800 \\ 136200 \\ 195000 \end{bmatrix}$

$$\text{d. } K_{38} = T^{38} \times K_0 = \begin{bmatrix} 0.80 & 0.09 & 0.10 \\ 0.12 & 0.76 & 0.05 \\ 0.08 & 0.15 & 0.85 \end{bmatrix}^{38} \begin{bmatrix} 300000 \\ 120000 \\ 180000 \end{bmatrix} = \begin{bmatrix} 194983 \\ 150513 \\ 254504 \end{bmatrix}$$

$$K_{50} = T^{50} \times K_0 = \begin{bmatrix} 0.80 & 0.09 & 0.10 \\ 0.12 & 0.76 & 0.05 \\ 0.08 & 0.15 & 0.85 \end{bmatrix}^{50} \begin{bmatrix} 300000 \\ 120000 \\ 180000 \end{bmatrix} = \begin{bmatrix} 194983 \\ 150513 \\ 254504 \end{bmatrix}$$

### Question 3

$$\text{a. } \begin{bmatrix} 2 & 1 & 1 \\ 1 & -1 & 1 \\ 0 & 2 & -1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 12 \\ 1 \\ 6 \end{bmatrix}$$

$$\text{b. Using a graphing calculator the determinant of the matrix } \begin{bmatrix} 2 & 1 & 1 \\ 1 & -1 & 1 \\ 0 & 2 & -1 \end{bmatrix} = \mathbf{A} \text{ is}$$

`det([A])`

`1`

■

So the equations have a unique solution because the determinant is non-zero.

c. Using a graphing calculator, the inverse matrix of A is

$$[A]^{-1} = \begin{bmatrix} -1 & 3 & 2 \\ 1 & -2 & -1 \\ 2 & -4 & -3 \end{bmatrix}$$

■

$$\mathbf{d.} \begin{bmatrix} -1 & 3 & 2 \\ 1 & -2 & 1 \\ 2 & -4 & -3 \end{bmatrix} \begin{bmatrix} 2 & 1 & 1 \\ 1 & -1 & 1 \\ 0 & 2 & -1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} -1 & 3 & 2 \\ 1 & -2 & 1 \\ 2 & -4 & -3 \end{bmatrix} \begin{bmatrix} 12 \\ 1 \\ 6 \end{bmatrix}$$

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 3 \\ 4 \\ 2 \end{bmatrix}$$

The ideal number is 3 bookshops, 4 sports shoe shops and 2 music stores.





© MATHEMATICAL ASSOCIATION OF VICTORIA 2007

Published by The Mathematical Association of Victoria

'Cliveden', 61 Blyth Street, Brunswick, 3056

Phone: (03) 9380 2399 Fax: (03) 9389 0399

E-mail: [office@mav.vic.edu.au](mailto:office@mav.vic.edu.au) website: <http://www.mav.vic.edu.au>

These solutions to the Victorian Curriculum and Assessment Authority 2006 VCE Mathematics Examination Papers have been written and published to assist students in their preparations for the Victorian Curriculum and Assessment Authority's 2007 VCE mathematics courses. These answers and solutions do not necessarily reflect the views of the Victorian Curriculum and Assessment Authority.

These solutions are licensed to the purchasing school or educational organisation with permission for copying within that school or educational organisation. No part of this publication may be reproduced, transmitted or distributed, in any form or by any means, outside purchasing schools or educational organisations or by individual purchasers, without permission.